

Topics in Primary Care Medicine

Osteoarthritis A Continuing Challenge

KENNETH E. SACK, MD, *San Francisco, California*

Osteoarthritis is a disorder of cartilage that affects almost 85% of the population by age 75. A lack of rigorous clinical and radiographic criteria for defining the disorder makes precise determination of its prevalence impossible. The process of wear and tear explains many manifestations of osteoarthritis, but it does not account for some of the clinical findings or the biochemical changes in osteoarthritic cartilage. Thus, other factors such as heredity, hormones, and diet may play a role. Treatment consists of teaching patients about their disease, alleviating pain, and preserving joint function. Nonsteroidal anti-inflammatory drugs may be no more effective than simple analgesics in relieving the pain of this disorder. Moreover, some nonsteroidal anti-inflammatory drugs can adversely affect cartilage metabolism, and most are possibly dangerous in elderly patients. Drugs that inhibit the production or activity of chondrolytic enzymes can slow the degeneration of cartilage in some animals, but their effects on humans with osteoarthritis are unproved. The surgical repair of severely damaged joints can have gratifying results.

(Sack KE: Osteoarthritis—A continuing challenge. *West J Med* 1995; 163:579-586)

Osteoarthritis is in many ways like the weather—ubiquitous, often unnoticed, sometimes dramatic in its effects, and largely unexplained. To carry the analogy further, everyone talks about osteoarthritis, but nobody seems to be doing much about it.

The name may be a misnomer. The disorder's primary cause is not likely to reside in bone,^{1,2} and inflammation is not prominent in its initial stages.³ Nevertheless, the term osteoarthritis serves its purpose—it is short, physicians and patients know what it means, and it has a more benign connotation for patients than does "degenerative joint disease."

The prevalence of osteoarthritis in the population is hard to determine because the disorder cannot be defined precisely. Do all patients with joint pain and radiographic evidence of osteophytes have osteoarthritis?^{4,5} Does a normal radiograph exclude its diagnosis in a patient with joint pain?⁶ Can osteoarthritis be completely asymptomatic?⁷ Are osteophytes a normal consequence of aging?⁴ Will our ability to quantify cartilaginous loss with magnetic resonance imaging scans alleviate or enhance our confusion?⁸

Osteoarthritis rarely occurs before age 40, but by age 75, at least 85% of the population have either clinical or radiographic evidence of the disease.⁹ Although it occurs more frequently and tends to affect more joints in women, for those younger than 45, it affects both sexes equally.¹⁰ White populations from developed countries have similar rates of osteoarthritis in the hands and the knees.¹⁰ By contrast, black American women have more osteoarthritis

of the knees.¹¹ Although it is relatively uncommon (less than 5%) in the hips in Chinese¹² and in most black populations,¹³ its prevalence in many European communities approaches 25%.¹³

What Causes Osteoarthritis?

For many years, investigators have embraced the concept that osteoarthritis is a disease of wear and tear. Some of its clinical manifestations do not fit this notion, however. For instance, why do women have a propensity for prominent bony enlargement of the distal interphalangeal (Heberden's nodes) and proximal interphalangeal (Bouchard's nodes) joints,¹⁴ whereas men who do manual labor show no such predisposition for the disease in their hands?^{15,16} Why does an inflammatory form of osteoarthritis tend to affect women near menopause?¹⁷⁻¹⁹ And why is cartilage in patients with the disorder biochemically different from cartilage in older people without it?²⁰

Understanding the structure and function of normal cartilage may provide some clues to the pathogenesis of osteoarthritis. Articular cartilage provides a low-friction bony interface greatly capable of absorbing shock. These functions are dependent on the composition of the cartilage matrix, which consists of collagen fibers (which make up 50% of the dry weight of cartilage) intermingled with proteoglycans—high-molecular-weight aggregates of glycosaminoglycans (mucopolysaccharides) bound to protein chains (Figure 1).²⁰⁻²² From a functional standpoint, collagen gives cartilage tensile strength and allows it to resist shear forces occurring during motion under




Figure 1.—A schematic representation of a proteoglycan aggregate is shown.

load. Collagen also constrains the negatively charged hydrated proteoglycans. This produces a large swelling pressure and gives cartilage its elasticity and resistance to compression.²¹ Under high-load conditions, proteoglycans release water and provide lubrication at the cartilage surface (hydrostatic lubrication).^{20,21} Not surprisingly, areas of cartilage that are subjected to heavy loads contain greater concentrations of proteoglycans.^{23,24}

Damage to the collagen fibers or to the proteoglycan matrix of cartilage causes a loss of its normal lubricating and shock-absorbing functions. Unfortunately, cartilage responds poorly to injury because of a limited blood supply and the lack of undifferentiated mesenchymal cells capable of initiating healing. Thus, repair tends to be spotty, and instead of containing collagen found in normal articular cartilage, the newly formed cartilage contains collagen characteristic of fibrocartilage.²⁵

The cartilage in patients with osteoarthritis is not simply “old and worn out” (Table 1). Compared with normal cartilage, aging cartilage contains less water, a higher ra-

TABLE 2.—Cytokines in Osteoarthritis

Cytokine	Effect
Interleukin (IL) 1	Stimulates loss of proteoglycan (PG) from cartilage, release of proteases, synthesis of IL-6 and type I collagen Inhibits cell division in chondrocytes, synthesis of type II collagen
Interferon gamma	Suppresses synthesis of type II collagen
Tumor necrosis factor alpha	Stimulates synthesis of proteases
IL-6	Inhibits IL-1-induced breakdown of cartilage and synthesis of cartilage PG Stimulates production of tissue inhibitor of metalloproteases (TIMP)
Transforming growth factor beta	Stimulates synthesis of collagen and TIMP Inhibits synthesis of collagenase
Insulin-like growth factor I ...	A potent anabolic factor of cartilage

tio of keratan to chondroitin sulfate, and fragmented link protein. Yet, there is no evidence that aging cartilage is biochemically inferior to normal cartilage. Cartilage in patients with osteoarthritis, on the other hand, contains more water, less keratan sulfate than in normal cartilage, and normal link protein.²⁰ Structural changes in the proteoglycan macromolecules are key features of osteoarthritis and, together with ultrastructural changes in collagen fibers, cause deterioration in the function of cartilage. The chondrocyte, previously thought to be an inert cell, is likely an important participant in this process. It has anabolic or catabolic functions, depending on local events. In contrast to rheumatoid arthritis where the synovium is the primary source of proteolytic enzymes, in osteoarthritis the chondrocyte seems to be the principal producer of these substances.^{26,27} Table 2 contains a partial list of cytokines likely to be important in the pathogenesis of osteoarthritis.^{27,28}

Role of Trauma

Could trauma initiate these biochemical events? Although loading and motion are important in preserving normal cartilage metabolism and function,²⁹⁻³³ exerting excessive force on cartilage can have the opposite effect.³⁴⁻³⁶ Thus, activities such as jumping or twisting can damage articular cartilage, and an injury that results in an unstable joint or that damages the normal cushioning structures, such as a meniscus in the knee, can render cartilage vulnerable to stress. It has been suggested that trauma to cartilage can be more subtle. Indeed, a subset of the normal population habitually loads their legs at heel strike and thus may be predisposing themselves to joint damage.³⁵ Having abnormally dense bone could also predispose cartilage to injury,³⁷⁻⁴¹ but this idea has been disputed.⁴²⁻⁴⁴ Finally, an intact neuromuscular system is important in maintaining normal joint function. Although deficits in sensory nerves are not an important cause of osteoarthritis, abnormalities in the muscular reflex that

TABLE 1.—Composition of Aging Cartilage Versus That Affected by Osteoarthritis*

Chemical Composition	Aging Cartilage	Osteoarthritic Cartilage
H ₂ O	↓	↑
Glycosaminoglycans	Normal or slightly ↓	↓
Keratan sulfate	↑	↓
Chondroitin sulfate	↓	Early ↑, then ↓
Hyaluronic acid	↑	↓
Proteoglycans	↑	‡
Link protein	Fragmented	Normal

↓ = amount decreased, ↑ = amount increased

*Modified from Brandt and Fife.²⁰
†Decreased extractability and normal aggregation.
‡Increased extractability and decreased aggregation.

normally protects the joints could increase its severity.⁴⁵

In 1971 a mechanism was proposed for the occurrence of “nodal” osteoarthritis in women.¹⁵ It was postulated that the smaller surface area of the distal joints of the fingers subjects these joints to greater pressures per square inch than the more proximal joints. Furthermore, fine motor activity uses the deep flexor tendons that insert into the distal phalanges, but these tendons are “splinted” by the superficial tendons during power-grip activities. Thus, knitting and sewing would likely produce more node formation than hammering and sawing. It was subsequently shown that in an industrial setting, node formation in the fingers correlated more with repetitive, fine motor activity than with gender.¹⁶

Role of Heredity

Early studies suggested that generalized osteoarthritis, which typically affects the hands, knees, and spine, occurs in two forms—one in association with Heberden’s nodes, the other without such nodes.⁴⁶⁻⁴⁸ Nodal osteoarthritis showed an inheritance pattern consistent with an autosomal gene, dominant in females and recessive in males.^{49,50} Non-nodal osteoarthritis showed a polygenic inheritance.⁵¹ Lacking in these studies, however, is information regarding occupation and other physical activities involving the hands. Current research has linked abnormalities in the type II procollagen gene *COL2A1* on chromosome 12 with some familial forms of generalized erosive osteoarthritis.^{52,53} Patients with these abnormalities typically have an associated chondrodystrophy, however. Such genetic defects do not account for most cases of osteoarthritis.⁵⁴

Role of Estrogen

The predisposition of women to severe osteoarthritis and the tendency for the inflammatory form to occur near menopause suggest that estrogens play a role in this disorder.⁵⁵ Indeed, the articular cartilage of some animals contains estrogen receptors.⁵⁶⁻⁵⁸ Whether estrogens alleviate or worsen the disease in animals seems to depend on the model being studied.^{59,60} Possibly detrimental effects of estrogens on cartilage include suppressing chondrocyte proliferation, suppressing proteoglycan synthesis, and stimulating cytokine release from macrophages and chondrocytes.⁶¹

Role of Diet

How might diet affect the development or severity of osteoarthritis? Certainly obesity plays a role in the occurrence and progression of the disease in the knees.⁶²⁻⁶⁶ This could be ascribed to the mechanical effect of increased weight, but it is hard to understand why obesity is a modest predictor of osteoarthritis in the distal interphalangeal and first carpometacarpal joints.⁶⁷ In some animal models, a diet high in saturated fat increases the severity of the disease,⁶⁸⁻⁷⁰ but in other studies this effect has been hard to reproduce.⁷¹ Some secondary forms of osteoarthritis have a presumed dietary association.⁷² Kashin-Beck disease,

for example, is a noninflammatory disorder of enchondral bone growth endemic to eastern Siberia, northern China, and northern Korea. Dystrophic changes in the epiphyseal and metaphyseal areas lead to secondary degenerative changes in peripheral joints and the spine. Proposed causes have included infectious agents and excesses or deficiencies of trace elements.⁷²⁻⁷⁴

Clinical Aspects of Osteoarthritis

Osteoarthritis affects most movable joints. Pain in a joint with movement along with articular bony enlargement are typical manifestations of, but are not specific for, the disease. Nevertheless, certain aspects of joint involvement are fairly consistent and deserve mention.

In the hands, osteoarthritis affects primarily the distal interphalangeal, proximal interphalangeal, and first carpometacarpal joints (Figure 2). Deformity or a decrease in motion of the joint occurs gradually, and deviation tends to be in a lateral direction, unlike the vertical deformity of rheumatoid arthritis. Metacarpophalangeal involvement is distinctly unusual, but elderly patients with extensive osteoarthritis of the hands occasionally have mild swelling or subluxation of the metacarpophalangeal joints. Heavy manual labor also may cause degenerative changes in these joints.⁷⁵

The wrists are virtually never involved, and if they are, this should suggest another diagnosis such as rheumatoid arthritis or pseudogout. Involvement of the first carpometacarpal joint, however, may give the wrist a “squared-off” appearance (see Figure 2).

Although osteoarthritis rarely affects the elbows in the absence of a metabolic disease or previous trauma, a subset of patients may have involvement of the elbows. These patients typically are men who have osteoarthritis in other joints, particularly the metacarpophalangeal joints.⁷⁶

Osteoarthritis frequently affects the acromioclavicular joint. By contrast, the glenohumeral joint is not a common target. When degenerative changes occur in this joint, patients are likely to have a severe tear in the rotator cuff, allowing superior migration of the humeral




Figure 2.—Bony enlargement is seen of the distal interphalangeal joint of the index finger (Heberden’s node) of a patient with osteoarthritis of the hand. The prominent first carpometacarpal joint gives the hand a squared-off appearance.

head.⁷⁷ Elderly women are susceptible to a destructive arthropathy of the shoulder (“Milwaukee shoulder”), presumably mediated by apatite crystals.⁷⁸

Movable portions of the spine are susceptible to degenerative changes. The central gelatinous portion of the intervertebral disc (nucleus pulposus) gradually disappears with aging and is essentially nonexistent after age 45.⁷⁹ Thus, “degeneration” of discs is a normal phenomenon of aging. In the lower five cervical vertebrae, such changes could place abnormal stress on the posterior (zygoapophyseal) vertebral articulations, resulting in joint-space narrowing, subarticular sclerosis, and osteophyte formation—the hallmarks of osteoarthritis. Because the cervical spine is reputed to move about 600 times in an hour (whether awake or asleep), it is no wonder that at any given time about 10% of the population has a pain in the neck.⁸⁰ Although radiographs of the cervical spine typically show bony abnormalities of the facet joints and the posterolateral uncinate processes, caution must be used in ascribing pain in the neck or radicular symptoms to such changes.⁸¹

Osteoarthritis affects the hips (often both) more commonly in men than in women.⁸² Heavy lifting might increase the risk of disease in this joint.⁸³ Other possible risk factors include leg-length discrepancy (the long leg will be affected), acetabular dysplasia, a slipped femoral epiphysis, and Legg-Calvé-Perthes disease. Most of the time, however, the cause is not apparent.⁸⁴ Pain from osteoarthritis in the hips typically occurs in the groin or anterior thigh, but occasionally in the buttock or even in the knees. Range of motion in the hip, particularly internal rotation, becomes limited. Weakness in the hip abductor muscles makes it difficult to keep the pelvis level during walking and causes a patient to tilt toward the affected side to swing the opposite leg forward (Trendelenburg gait).

The prevalence of osteoarthritis in the knees rises with age.^{65,82,85} Although a third of people older than 65 have radiographic findings in the knee consistent with the disease, only 10% of this age group has pain in the knees and an abnormality of cartilage by visual or radiographic examination.^{65,85} Obesity correlates strongly with osteoarthritis of the knees, particularly in women and in patients with disease of both knees.⁶⁴ Obese patients and those with a previous meniscectomy have more involvement of the tibiofemoral than the patellofemoral compartment.⁶⁶ Isolated patellofemoral disease is more common in young adults, either as a result of trauma or, in women, from recurrent patellar subluxation.⁸⁶ Although habitual physical activity of many types, including running, does not predispose to disease in the knees,^{87,88} prolonged or repeated kneeling and squatting appear to be risk factors.³⁶ Characteristic physical findings include pain or crepitation on joint motion, tenderness along the tibiofemoral or patellofemoral articulations, and genu valgum (knock-knee) or genu varum (bowleg) deformity.

In the absence of repetitive trauma or chronic ligamentous injury, osteoarthritis seldom affects the an-

TABLE 3.—Other Conditions Causing Degeneration of Cartilage*

Cause	Condition
Inflammatory diseases	Rheumatoid arthritis Infection
Physical factors	Trauma—physical injury, obesity, mechanical derangement, neuropathic joint Ischemic necrosis of bone
Endocrine diseases	Diabetes mellitus Acromegaly
Metabolic diseases	Hemochromatosis Wilson's disease Ochronosis Crystalline—urate, calcium pyrophosphate, hydroxyapatite arthropathy
Nutritional (?)	Kashin-Beck disease Steroid injections (?)

*Modified from Sack.⁹⁷

kles.^{82,87,89} The first metatarsophalangeal joint, however, is a common target. Inflammation of the bursa overlying the medial aspect of this joint may produce a characteristic swelling (bunion).

Looking at the pattern of osteoarthritis, the impression is that humans may have evolved faster than their joints could adapt.^{90,91} After all, we developed an opposable thumb, assumed an upright posture, and learned to live well beyond our reproductive years. And we cannot regenerate new limbs.

Variations of Osteoarthritis

A “nodal generalized” form of osteoarthritis affects principally the distal and proximal interphalangeal and first carpometacarpal joints of the hands, but it also involves the hips, knees, metatarsophalangeal joints, and the spinal articulations.^{46,48} Joint inflammation and exaggerated osteophyte formation can occur. An “erosive” type of osteoarthritis shows many features of generalized disease, but joint inflammation and subchondral erosive changes are more pronounced.⁹²⁻⁹⁴ Patients with this form of the disease are typically women who have recently entered menopause.^{55,95} The female predilection for nodal generalized osteoarthritis and the reported association of this disorder with the HLA-A1, B8 tissue type, Sjögren's syndrome, and thyroid disease suggest that this is an autoimmune disorder.⁹⁶

When osteoarthritis occurs at atypical sites or is unusually severe, other factors may be operative (Table 3).⁹⁷ As a rule, any physical, inflammatory, or metabolic insult can predispose cartilage to further degenerative change. Particularly noteworthy is the occurrence of severe osteoarthritis in association with the deposition of calcium pyrophosphate dihydrate or apatite crystals.^{98,99}

Laboratory Evaluation

Osteoarthritis causes few, if any, abnormal laboratory values. The erythrocyte sedimentation rate is typically normal but may show a modest increase in patients with

inflammatory osteoarthritis. Tests for rheumatoid factor or antinuclear antibodies are often positive in normal aging persons and therefore have little meaning when the clinical findings suggest only osteoarthritis. Synovial fluid obtained from joints affected by the disease typically shows normal viscosity and few leukocytes ($<2 \times 10^9$ per liter [2,000 per mm³]) with a normal differential count (<0.3 [30%] polymorphonuclear cells). In rare cases, fluid from joints that have severe degeneration of cartilage or an associated deposition of crystals (such as calcium pyrophosphate dihydrate or apatite) will show a higher leukocyte count.

Radiographs in patients with osteoarthritis may show narrowing of the joint space, a marginal overgrowth of bone (osteophytes), subchondral bony sclerosis or cyst formation, or malalignment of the joint. Although these findings frequently constitute criteria for the diagnosis, they do not always correlate with clinical symptoms or with the condition of articular cartilage on arthroscopy.^{100,101} Some radiographic changes, however, have prognostic importance. Thus, in hips, supralateral migration of the femoral head may carry a worse prognosis than medial migration.¹⁰² Magnetic resonance imaging may show degenerative changes in cartilage when the plain radiograph does not.¹⁰³ Such studies are costly, however, and are useful only when planning surgical intervention.

Treatment

Grandma Moses treated her arthritis with sweet milk and turpentine.¹⁰⁴ The question is, can we do any better now? If the goal is to repair damaged cartilage, the answer is clearly no. If, however, the goal is to teach patients about their disease, alleviate pain, preserve joint function, or ultimately repair a severely damaged joint, then the answer is yes.

We can now provide patients a clearer picture of their long-term prognosis. For instance, studies show that osteoarthritis seldom impairs function of the hand, even if some pain and stiffness persist.^{105,106} Furthermore, pain in the knees or hips may diminish as patients reach their mid-60s, even if damage to these joints is substantial.^{107,108} And never underestimate the power of reassurance. In one study, patients who had monthly phone calls from lay personnel to discuss self-care issues had a substantial reduction of joint pain.¹⁰⁹

Physical modalities provide the cornerstone for managing osteoarthritis. Activities that involve normal loading of joints are important in maintaining healthy cartilage.²⁹⁻³³ A knowledgeable physical or occupational therapist can teach patients how to exercise safely and how to protect their joints while doing routine activities of daily living. For patients with disease of the hips, the simple measure of splitting heavy loads and carrying them in each hand, or carrying the load ipsilateral to the affected hip, will minimize the force exerted on that joint.¹¹⁰ In osteoarthritis of the knees, weakness of the quadriceps muscle¹¹¹ and obesity correlate with symptoms.⁶²⁻⁶⁷ Thus, sensible treatment would include muscle strengthening¹¹² and weight reduction.¹¹³ Modifying shoes will occasion-

ally improve the mechanics in knees¹¹⁴ and relieve pain. Although it is hard to prove that physical modalities alter the course of osteoarthritis, such intervention may at least lessen symptoms. Two recent articles review the principles of exercise for patients with osteoarthritis.^{30,115}

The cause of pain in this disease is conjectural.^{116,117} Cartilage contains no nerve fibers, and synovitis does not always correlate with symptoms. Thus, it is no surprise that NSAIDs may be no better than simple analgesics like acetaminophen at relieving the pain of osteoarthritis.¹¹⁸⁻¹²¹ Some NSAIDs adversely affect proteoglycan metabolism and could theoretically inhibit repair mechanisms in cartilage, but the effect of any particular NSAID on the eventual outcome of osteoarthritis is unknown.¹²²⁻¹²⁷ Because these drugs are relatively toxic to elderly patients—the population at risk for having osteoarthritis—it is prudent to use them in their lowest effective dosage and only if they relieve symptoms more than simple analgesics.

Administering steroids intra-articularly can reduce pain in joints affected by arthritis.¹²⁸ This would indicate that inflammation has a role in this process, but placebo responses could also account for some of the “successes” of intra-articular steroid use in this disorder. Intra-articularly or parenterally administered steroids could affect chondrocyte metabolism and the production of cytokines and proteolytic enzymes, but studies of the effects of such treatment on experimental osteoarthritis give conflicting results.¹²⁹⁻¹³⁸ Conceivably intra-articular steroid use could hasten the progression of the disease, but at least one study indicates that it takes less steroid to suppress chondrocyte protease synthesis than to suppress proteoglycan synthesis.¹³⁸

Chloroquine and doxycycline can inhibit the production or activity of chondrolytic enzymes.¹³⁹ Calcitonin has anabolic effects in chondrocytes and can stimulate cartilage growth.²⁵ Although these substances have shown some promise in experimental models, it remains to be seen whether they will benefit patients with established osteoarthritis. Unfortunately, the positive results from the intra-articular administration of a chondroprotective agent such as hyaluronan¹⁴¹⁻¹⁴³ could not be duplicated in a recent controlled trial.¹⁴⁴ Applied topically, capsaicin, which depletes local sensory nerve terminals of substance P, can reduce pain and tenderness in small joints affected by osteoarthritis.^{145,146} The long-term effects of such therapy are unknown.

Most patients with osteoarthritis will never need surgical treatment of their joints, but when pain is unrelenting or when joint function is severely compromised, such treatment can provide substantial benefit. The most gratifying results have come from remodeling or replacing a hip or knee. Surgical therapy is rarely necessary for osteoarthritis of the hand, but occasionally a patient may benefit from fusing, resecting, or replacing a joint. Similar procedures are helpful when the disease causes severe pain or deformity in the feet.

Arthroscopic debridement—trimming meniscal fragments, removing loose debris, shaving shaggy articular surfaces—may relieve symptoms in some patients with

disease of the knee. Abrading the articular surface does not offer any additional benefit.¹⁴⁷ Some investigators advocate lavaging the knee by means of an arthroscope or cannula in an attempt to remove phlogistic debris.¹⁴⁸ The long-term results of such treatment are unknown.

The Challenge

To meet the challenge of this disease, we must clarify the factors leading to its onset and progression and devise safe and practical methods of preserving cartilage and relieving pain. The recent report that cultured autologous chondrocytes can be used to repair deep cartilage defects is a step in the right direction.¹⁴⁹

REFERENCES

- Price T, Hesp R, Mitchell R: Bone density in generalized osteoarthritis. *J Rheumatol* 1987; 14:560-562
- Knight SM, Ring EF, Bhalla AK: Bone mineral density and osteoarthritis. *Ann Rheum Dis* 1992; 51:1025-1026
- Howell DS, Sapolsky AI, Pita JC, Woessner F: The pathogenesis of osteoarthritis. *Semin Arthritis Rheum* 1976; 4:365-383
- Hernborg J, Nilsson BE: The relationship between osteophytes in the knee joint, osteoarthritis and aging. *Acta Orthop Scand* 1973; 44:69-74
- Hadler NM: Knee pain is the malady—Not osteoarthritis. *Ann Intern Med* 1992; 116:598-599
- Kaye JJ: Radiologic assessment of osteoarthritis—New techniques. *Rheum Dis Clin North Am* 1993; 19:659-672
- Davis MA, Ettinger WH, Neuhaus JM, Barclay JD, Segal MR: Correlates of knee pain among US adults with and without radiographic knee osteoarthritis. *J Rheumatol* 1992; 19:1943-1949
- Peterfy CG, Majumdar S, Lang P, van Dijke CF, Sack K, Genant HK: MR imaging of the arthritic knee: Improved discrimination of cartilage, synovium, and effusion with pulsed saturation transfer and fat-suppressed T1-weighted sequences. *Radiology* 1994; 191:413-419
- Lawrence JS, Bremner JM, Bier F: Osteoarthritis—Prevalence in the population and relationship between symptoms and x-ray changes. *Ann Rheum Dis* 1966; 25:1-24
- Felson DT: Osteoarthritis. *Rheum Dis Clin North Am* 1990; 16:499-512
- Anderson JJ, Felson DT: Factors associated with osteoarthritis of the knee in the first national Health and Nutrition Examination Survey (HANES I)—Evidence for an association with overweight, race, and physical demands of work. *Am J Epidemiol* 1988; 128:179-189
- Hoaglund FT, Yau AC, Wong WL: Osteoarthritis of the hip and other joints in southern Chinese in Hong Kong. *J Bone Joint Surg (Am)* 1973; 55:645-657
- Lawrence J, Sebo M: The geography of osteoarthritis. In Nuki G (Ed): *The Aetio-pathogenesis of Osteoarthritis*. Baltimore, Md, University Park Press, 1958, pp 155-183
- Stecher R: Heberden's nodes—A disease description of osteoarthritis in the finger joint. *Ann Rheum Dis* 1955; 14:1-10
- Radin EL, Parker HG, Paul IL: Pattern of degenerative arthritis—Preferential involvement of distal finger-joints. *Lancet* 1971; 1:377-379
- Hadler NM, Gillings DB, Imbus HR, et al: Hand structure and function in an industrial setting. *Arthritis Rheum* 1978; 21:210-220
- Crain D: Interphalangeal osteoarthritis. *JAMA* 1961; 175:1049-1053
- Ehrlich GE: Inflammatory osteoarthritis—I. The clinical syndrome. *J Chronic Dis* 1972; 25:317-328
- Belhorn LR, Hess EV: Erosive osteoarthritis. *Semin Arthritis Rheum* 1993; 22:298-306
- Brandt KD, Fife RS: Ageing in relation to the pathogenesis of osteoarthritis. *Clin Rheum Dis* 1986; 12:117-130
- Greenwald RA, Moy WW, Seibold J: Functional properties of cartilage proteoglycans. *Semin Arthritis Rheum* 1978; 8:53-67
- Hardingham TE, Bayliss M: Proteoglycans of articular cartilage: Changes in aging and in joint disease. *Semin Arthritis Rheum* 1990; 20(suppl 1):12-33
- Yoshida K, Azuma H: Contents and compositions of glycosaminoglycans in different sites of the human hip joint cartilage. *Ann Rheum Dis* 1982; 41:512-519
- Kiviranta I, Jurvelin J, Tammi M, Säämänen AM, Helminen HJ: Weight bearing controls glycosaminoglycan concentration and articular cartilage thickness in the knee joints of young beagle dogs. *Arthritis Rheum* 1987; 30:801-809
- Howell DS, Altman RD: Cartilage repair and conservation in osteoarthritis—A brief review of some experimental approaches to chondroprotection. *Rheum Dis Clin North Am* 1993; 19:712-724
- Poole A: Cartilage in health and disease. In McCarty D, Koopman W (Eds): *Arthritis and Allied Conditions—A Textbook of Rheumatology*. Philadelphia, Pa, Lea & Febiger, 1993, pp 279-333
- Pelletier JP, DiBattista JA, Roughley P, McCollum R, Martel-Pelletier J: Cytokines and inflammation in cartilage degradation. *Rheum Dis Clin North Am* 1993; 19:545-568
- Pelletier JP, Roughley PJ, DiBattista JA, McCollum R, Martel-Pelletier J: Are cytokines involved in osteoarthritic pathophysiology? *Semin Arthritis Rheum* 1991; 20(suppl 2):12-25
- Palmoski MJ, Colyer RA, Brandt KD: Joint motion in the absence of normal loading does not maintain normal articular cartilage. *Arthritis Rheum* 1980; 23:325-334
- Allen ME: Arthritis and adaptive walking and running. *Rheum Dis Clin North Am* 1990; 16:887-914
- Radin EL, Burr DB: Hypothesis: Joints can heal. *Semin Arthritis Rheum* 1984; 13:293-302
- Palmoski M, Perricone E, Brandt KD: Development and reversal of a proteoglycan aggregation defect in normal canine knee cartilage after immobilization. *Arthritis Rheum* 1979; 22:508-517
- Salter RB, Simmonds DF, Malcolm BW, Rumble EJ, MacMichael D, Clements ND: The biological effect of continuous passive motion on the healing of full-thickness defects in articular cartilage. *J Bone Joint Surg (Am)* 1980; 62:1232-1251
- Radin EL, Paul IL: Response of joints to impact loading—I. In vitro wear. *Arthritis Rheum* 1971; 14:356-362
- Radin EL, Burr DB, Caterson B, Fyhrrie D, Brown TD, Boyd RD: Mechanical determinants of osteoarthritis. *Semin Arthritis Rheum* 1991; 21(suppl 2):12-21
- Cooper C, McAlindon T, Coggon D, Egger P, Dieppe P: Occupational activity and osteoarthritis of the knee. *Ann Rheum Dis* 1994; 53:90-93
- Belmonte-Serrano MA, Bloch DA, Lane NE, Michel BE, Fries JF: The relationship between spinal and peripheral osteoarthritis and bone density measurements. *J Rheumatol* 1993; 20:1005-1013
- Hannan MT, Anderson JJ, Zhang Y, Levy D, Felson DT: Bone mineral density and knee osteoarthritis in elderly men and women—The Framingham Study. *Arthritis Rheum* 1993; 36:1671-1680
- Radin EL, Swann DA, Paul IL, McGrath PJ: Factors influencing articular cartilage wear in vitro. *Arthritis Rheum* 1982; 25:974-980
- Cumming RG, Klineberg RJ: Epidemiological study of the relation between arthritis of the hip and hip fractures. *Ann Rheum Dis* 1993; 52:707-710
- Dequeker J: The relationship between osteoporosis and osteoarthritis. *Clin Rheum Dis* 1985; 11:271-296
- Dedrick DK, Goldstein SA, Brandt KD, O'Connor BL, Goulet RW, Albrecht M: A longitudinal study of subchondral plate and trabecular bone in cruciate-deficient dogs with osteoarthritis followed up for 54 months. *Arthritis Rheum* 1993; 36:1460-1467
- Hordon LD, Wright V, Smith MA: Bone mass in osteoarthritis. *Ann Rheum Dis* 1992; 51:823-825
- Hochberg M, Lethbridge-Cejku M, Scott W Jr, Plato C, Tobin J: Appendicular bone mass and osteoarthritis of the hands in women: Data from the Baltimore Longitudinal Study of Aging. *J Rheumatol* 1994; 21:1532-1536
- O'Connor BL, Brandt KD: Neurogenic factors in the etiopathogenesis of osteoarthritis. *Rheum Dis Clin North Am* 1993; 19:581-605
- Kellgren J, Moore R: Generalized osteoarthritis and Heberden's nodes. *BMJ* 1952; 1:181-187
- Kellgren J, Lawrence J: Osteoarthrosis and disk degeneration in an urban population. *Ann Rheum Dis* 1958; 17:388-397
- Lawrence J: Generalized osteoarthrosis in a population sample. *Am J Epidemiol* 1960; 90:381-389
- Stecher R: Heberden oration: Heberden's nodes—A clinical description of osteoarthrosis of the finger joints. *Ann Rheum Dis* 1955; 14:1-10
- Stecher R, Hersh A, Hauser H: Heberden's nodes—The family history and radiographic appearance of a large family. *Am J Hum Genet* 1953; 5:46-60
- Lawrence J, Gelsthorpe K, Morell G: Heberden's nodes and HLA markers in generalized osteoarthritis. *J Rheumatol* 1983; 10(suppl 9):32-33
- Katzenstein PL, Maleski CJ, Pathria MN, Carter JR, Sheon RP, Moskowitz RW: Early-onset primary osteoarthritis and mild chondrodysplasia: Radiographic and pathologic studies with an analysis of cartilage proteoglycans. *Arthritis Rheum* 1990; 33:674-684
- Ala-Kokko L, Baldwin CT, Moskowitz RW, Prockop DJ: Single base mutation in the type II procollagen gene (COL2A1) as a cause of primary osteoarthritis associated with a mild chondrodysplasia. *Proc Natl Acad Sci USA* 1990; 87:6565-6568
- Vikkula M, Nissilä M, Hirvensalo E, et al: Multiallelic polymorphism of the cartilage collagen gene: No association with osteoarthrosis. *Ann Rheum Dis* 1993; 52:762-764
- Rosner IA, Goldberg VM, Moskowitz RW: Estrogens and osteoarthritis. *Clin Orthop* 1986; (213):77-83

56. Young PC, Stack MT: Estrogen and glucocorticoid receptors in adult canine articular cartilage. *Arthritis Rheum* 1982; 25:568-573
57. Rosner IA, Manni A, Malemud CJ, Boja B, Moskowitz RW: Estradiol receptors in articular chondrocytes. *Biochem Biophys Res Commun* 1982; 106:1378-1382
58. Sheridan P, Aufdemorte T, Holt G, Gates G: Cartilage of the baboon contains estrogen receptors. *Rheum Int* 1985; 5:279-281
59. Silberberg M, Silberberg R: Modifying action of estrogen on the evaluation of osteoarthritis in mice of different ages. *J Endocrinol* 1963; 72:449-451
60. Rosner IA, Goldberg VM, Getzy L, Moskowitz RW: Effects of estrogen on cartilage and experimentally induced osteoarthritis. *Arthritis Rheum* 1979; 22:52-58
61. Chander CL, Spector TD: Oestrogens, joint disease and cartilage. *Ann Rheum Dis* 1991; 50:139-140
62. Felson DT, Anderson JJ, Naimark A, Walker AM, Meenan RF: Obesity and knee osteoarthritis—The Framingham Study. *Ann Intern Med* 1988; 109:18-24
63. Kohatsu ND, Schurman DJ: Risk factors for the development of osteoarthritis of the knee. *Clin Orthop* 1990; (261):242-246
64. Davis MA, Ettinger WH, Neuhaus JM: Obesity and osteoarthritis of the knee: Evidence from the National Health and Nutrition Examination Survey (NHANES I). *Semin Arthritis Rheum* 1990; 20(suppl 1):34-41
65. Felson DT: The epidemiology of knee osteoarthritis: Results from the Framingham Osteoarthritis Study. *Semin Arthritis Rheum* 1990; 20(suppl 1):42-50
66. Cooper C, McAlindon T, Snow S, et al: Mechanical and constitutional risk factors for symptomatic knee osteoarthritis: Differences between medial tibiofemoral and patellofemoral disease. *J Rheumatol* 1994; 21:307-313
67. Hart DJ, Spector TD: The relationship of obesity, fat distribution and osteoarthritis in women in the general population—The Chingford Study. *J Rheumatol* 1993; 20:331-335
68. Silberberg M, Silberberg R: Effects of a high fat diet on the joints of aging mice. *Arch Pathol* 1950; 50:828-846
69. Silberberg M, Jarrett S, Silberberg R: Obesity and degenerative joint disease. *Arch Pathol* 1956; 61:280-288
70. Silberberg M, Silberberg R: Osteoarthritis in mice fed diets enriched with animal or vegetable fat. *Arch Pathol* 1960; 70:381-390
71. Sokoloff L, Mickelsen O: Dietary fat supplements, body weight and osteoarthritis in DBA/2JN mice. *Nutrition* 1965; 85:117-121
72. Sokoloff L: Endemic forms of osteoarthritis. *Clin Rheum Dis* 1985; 11:187-202
73. Sokoloff L: Kashin-Beck disease. *Rheum Dis Clin North Am* 1987; 13:101-104
74. Zhang GQ, Liu JX: An experimental animal model of Kashin-Beck disease. *Ann Rheum Dis* 1989; 48:149-152
75. Williams WV, Cope R, Gaunt WD, et al: Metacarpophalangeal arthropathy associated with manual labor (Missouri metacarpal syndrome)—Clinical, radiographic, and pathologic characteristics of an unusual degeneration process. *Arthritis Rheum* 1987; 30:1362-1371
76. Doherty M, Preston B: Primary osteoarthritis of the elbow. *Ann Rheum Dis* 1989; 48:743-747
77. Neer CS 2d, Craig EV, Fukuda H: Cuff-tear arthropathy. *J Bone Joint Surg (Am)* 1983; 65:1232-1244
78. McCarty DJ, Halverson PB, Carrera GF, Brewer BJ, Kozin F: 'Milwaukee shoulder'—Association of microspideroids containing hydroxyapatite crystals, active collagenase, and neutral protease with rotator cuff defects—I. Clinical aspects. *Arthritis Rheum* 1981; 24:464-473
79. Bland JH: Diagnosis and treatment of osteoarthritis. *Curr Opin Rheumatol* 1989; 1:315-324
80. Bland JH, Boushey DR: Anatomy and physiology of the cervical spine. *Semin Arthritis Rheum* 1990; 20:1-20
81. Harfing J, Halla J: Cervical spine syndromes. In McCarty D, Koopman W (Eds): *Arthritis and Allied Conditions—A Textbook of Rheumatology*. Philadelphia, Pa, Lea & Febiger, 1993, pp 1563-1571
82. Cushnaghan J, Dieppe P: Study of 50 patients with limb joint osteoarthritis—I. Analysis by age, sex, and distribution of symptomatic joint sites. *Ann Rheum Dis* 1991; 50:8-13
83. Croft P, Coggon D, Cruddas M, Cooper C: Osteoarthritis of the hip: An occupational disease in farmers. *BMJ* 1992; 304:1269-1272
84. Hoaglund FT, Shiba R, Newberg AH, Leung KY: Diseases of the hip. *J Bone Joint Surg (Am)* 1985; 67:1376-1383
85. Felson DT, Naimark A, Anderson JJ, Kazis L, Castelli W, Meenan RF: The prevalence of knee osteoarthritis in the elderly—The Framingham Osteoarthritis Study. *Arthritis Rheum* 1987; 30:914-918
86. Felson DT, Radin EL: What causes knee osteoarthritis: Are different compartments susceptible to different risk factors? *J Rheumatol* 1994; 21:181-183
87. Lane NE, Buckwalter JA: Exercise: A cause of osteoarthritis? *Rheum Dis Clin North Am* 1993; 19:617-633
88. Hannan MT, Felson DT, Anderson JJ, Naimark A: Habitual physical activity is not associated with knee osteoarthritis: The Framingham Study. *J Rheumatol* 1993; 20:704-709
89. Panush RS: Does exercise cause arthritis?—Long-term consequences of exercise on the musculoskeletal system. *Rheum Dis Clin North Am* 1990; 16:827-836
90. Hutton CW: Generalised osteoarthritis: An evolutionary problem? *Lancet* 1987; 1:1463-1465
91. Hutton CW: Osteoarthritis: The cause not result of joint failure? *Ann Rheum Dis* 1989; 48:958-961
92. Peter JB, Pearson CM, Marmor L: Erosive osteoarthritis of the hands. *Arthritis Rheum* 1966; 9:365-388
93. Cobby M, Cushnaghan J, Creamer P, Dieppe P, Watt I: Erosive osteoarthritis: Is it a separate disease entity? *Clin Radiol* 1990; 42:258-263
94. Altman RD, Gray R: Inflammation in osteoarthritis. *Clin Rheum Dis* 1985; 11:353-365
95. Ehrlich GE: Osteoarthritis beginning with inflammation—Definitions and correlations. *JAMA* 1975; 232:157-159
96. Doherty M, Patrick M, Powell R: Nodal generalised osteoarthritis is an autoimmune disease. *Ann Rheum Dis* 1990; 49:1017-1020
97. Sack KE: Osteoarthritis: Cause and long-term management. *Compr Ther* 1982; 8:46-51
98. Gibilisco PA, Schumacher HR Jr, Hollander JL, Soper KA: Synovial fluid crystals in osteoarthritis. *Arthritis Rheum* 1985; 28:511-515
99. Schumacher HR Jr: Crystals, inflammation, and osteoarthritis. *Am J Med* 1987; 83(5A):11-16
100. Blackburn WD Jr, Bernreuter WK, Rominger M, Loose LL: Arthroscopic evaluation of knee articular cartilage: A comparison with plain radiographs and magnetic resonance imaging. *J Rheumatol* 1994; 21:675-679
101. Fife RS, Brandt KD, Braunstein EM, et al: Relationship between arthroscopic evidence of cartilage damage and radiographic evidence of joint space narrowing in early osteoarthritis of the knee. *Arthritis Rheum* 1991; 34:377-382
102. Ledingham J, Dawson S, Preston B, Milligan G, Doherty M: Radiographic progression of hospital referred osteoarthritis of the hip. *Ann Rheum Dis* 1993; 52:263-267
103. Chan WP, Lang P, Stevens MP, et al: Osteoarthritis of the knee: Comparison of radiography, CT, and MR imaging to assess extent and severity. *AJR Am J Roentgenol* 1991; 157:799-806
104. Kallir O: *Grandma Moses—My Life's History*. New York, NY, Harper & Brothers, 1952
105. Baron M, Dutil E, Berkson L, Lander P, Becker R: Hand function in the elderly: Relation to osteoarthritis. *J Rheumatol* 1987; 14:815-819
106. Patrick M, Aldridge S, Hamilton E, Manhire A, Doherty M: A controlled study of hand function in nodal and erosive osteoarthritis. *Ann Rheum Dis* 1989; 48:978-982
107. Forman MD, Malament R, Kaplan D: A survey of osteoarthritis of the knee in the elderly. *J Rheumatol* 1983; 10:282-287
108. Felson DT: The course of osteoarthritis and factors that affect it. *Rheum Dis Clin North Am* 1993; 19:607-615
109. René J, Weinberger M, Mazzuca SA, Brandt KD, Katz BP: Reduction of joint pain in patients with knee osteoarthritis who have received monthly telephone calls from lay personnel and whose medical treatment regimens have remained stable. *Arthritis Rheum* 1992; 35:511-515
110. Neumann DA, Cook TM, Sholly RL, Sobush DC: An electromyographic analysis of hip abductor muscle activity when subjects are carrying loads in one or both hands. *Phys Ther* 1992; 72:207-217
111. McAlindon TE, Cooper C, Kirwan JR, Dieppe PA: Determinants of disability in osteoarthritis of the knee. *Ann Rheum Dis* 1993; 52:258-262
112. Fisher NM, Pendergast DR, Gresham GE, Calkins E: Muscle rehabilitation: Its effect on muscular and functional performance of patients with knee osteoarthritis. *Arch Phys Med Rehabil* 1991; 72:367-374
113. Felson DT, Zhang Y, Anthony JM, Naimark A, Anderson JJ: Weight loss reduces the risk for symptomatic knee osteoarthritis in women—The Framingham Study. *Ann Intern Med* 1992; 116:535-539
114. Yasuda K, Sasaki T: The mechanics of treatment of the osteoarthritic knee with a wedged insole. *Clin Orthop* 1987; (215):162-172
115. Bunning RD, Materson RS: A rational program of exercise for patients with osteoarthritis. *Semin Arthritis Rheum* 1991; 21 (suppl 2):33-43
116. Hadler NM: Osteoarthritis as a public health problem. *Clin Rheum Dis* 1985; 11:175-185
117. Altman RD, Dean D: Pain in osteoarthritis—Introduction and overview. *Semin Arthritis Rheum* 1989; 18(suppl 2):1-3
118. Bradley JD, Brandt KD, Katz BP, Kalasinski LA, Ryan SI: Comparison of an antiinflammatory dose of ibuprofen, an analgesic dose of ibuprofen, and acetaminophen in the treatment of patients with osteoarthritis of the knee. *N Engl J Med* 1991; 325:87-91
119. Bradley JD, Brandt KD, Katz BP, Kalasinski LA, Ryan SI: Treatment of knee osteoarthritis: Relationship of clinical features of joint inflammation to the

- response to a nonsteroidal antiinflammatory drug or pure analgesic. *J Rheumatol* 1992; 19:1950-1954
120. Dieppe P, Cushnaghan J, Jasani MK, McCrae F, Wan I: A two-year, placebo-controlled trial of non-steroidal anti-inflammatory therapy in osteoarthritis of the knee joint. *Br J Rheumatol* 1993; 32:595-600
 121. Williams HJ, Ward JR, Egger MJ, et al: Comparison of naproxen and acetaminophen in a two-year study of treatment of osteoarthritis of the knee. *Arthritis Rheum* 1993; 36:1196-1206
 122. Palmoski MJ, Colyer RA, Brandt KD: Marked suppression by salicylate of the augmented proteoglycan synthesis in osteoarthritic cartilage. *Arthritis Rheum* 1980; 23:83-91
 123. Palmoski MJ, Brandt KD: Effects of some nonsteroidal antiinflammatory drugs on proteoglycan metabolism and organization in canine articular cartilage. *Arthritis Rheum* 1980; 23:1010-1020
 124. Palmoski MJ, Brandt KD: Relationship between matrix proteoglycan content and the effects of salicylate and indomethacin on articular cartilage. *Arthritis Rheum* 1983; 26:528-531
 125. Herman JH, Hess EV: Nonsteroidal anti-inflammatory drugs and modulation of cartilaginous changes in osteoarthritis and rheumatoid arthritis—Clinical implications. *Am J Med* 1984; 77(4B):16-25
 126. Brandt KD: Effects of nonsteroidal anti-inflammatory drugs on chondrocyte metabolism in vitro and in vivo. *Am J Med* 1987; 83(5A):29-34
 127. Brandt KD: Should nonsteroidal anti-inflammatory drugs be used to treat osteoarthritis? *Rheum Dis Clin North Am* 1993; 19:29-44
 128. Hollinder J: Osteoarthritis: Perspectives on treatment. *Postgrad Med* 1980; 68:161-168
 129. Behrens F, Shepard N, Mitchell N: Alterations of rabbit articular cartilage by intra-articular injections of glucocorticoids. *J Bone Joint Surg (Am)* 1975; 57:70-76
 130. Behrens F, Shepard N, Mitchell N: Metabolic recovery of articular cartilage after intra-articular injections of glucocorticoid. *J Bone Joint Surg (Am)* 1976; 58:1157-1160
 131. Silberman M, von der Mark K, Maor G, van Menxel M: Dexamethasone impairs growth and collagen synthesis in condylar cartilage in vitro. *Bone Miner* 1967; 2:87-98
 132. Myers SL, Brandt KD, O'Connor BL: Low dose prednisone treatment does not reduce the severity of osteoarthritis in dogs after anterior cruciate ligament transection. *J Rheumatol* 1991; 18:1856-1862
 133. Butler M, Colombo C, Hickman L, et al: A new model of osteoarthritis in rabbits—Evaluation of anti-osteoarthritic effects of selected drugs administered intraarticularly. *Arthritis Rheum* 1983; 26:1380-1386
 134. Pelletier JP, Martel-Pelletier J, Cloutier JM, Woessner JF Jr: Proteoglycan-degrading acid metalloprotease activity in human osteoarthritic cartilage, and the effect of intraarticular steroid injections. *Arthritis Rheum* 1987; 30:541-548
 135. Pelletier JP, Martel-Pelletier J: Protective effects of corticosteroids on cartilage lesions and osteophyte formation in the Pond-Nuki dog model of osteoarthritis. *Arthritis Rheum* 1989; 32:181-193
 136. van der Kraan PM, Vitters EL, Postma NS, Verbunt J, van den Berg WB: Maintenance of the synthesis of large proteoglycans in anatomically intact murine articular cartilage by steroids and insulin-like growth factor I. *Ann Rheum Dis* 1993; 52:734-741
 137. Pelletier JP, Mineau F, Raynauld JP, Woessner JF Jr, Gunja-Smith Z, Martel-Pelletier JP: Intraarticular injections with methylprednisolone acetate reduce osteoarthritic lesions in parallel with chondrocyte stromelysin synthesis in experimental osteoarthritis. *Arthritis Rheum* 1994; 37:414-423
 138. Pelletier JP, Cloutier JM, Martel-Pelletier J: In vitro effects of tiaprofenic acid, sodium salicylate and hydrocortisone on the proteoglycan metabolism of human osteoarthritic cartilage. *J Rheumatol* 1989; 16:646-655
 139. Fife R, Brandt KD: Other approaches to therapy. In Moskowitz R, Howell D, Goldberg V, Mankin H (Eds): *Osteoarthritis: Diagnosis and Medical/Surgical Management*, 2nd edition. Philadelphia, Pa. WB Saunders, 1992, pp 511-526
 140. Yu LP Jr, Smith GN Jr, Brandt KD, Myers SL, O'Connor BL, Brandt DA: Reduction of the severity of canine osteoarthritis by prophylactic treatment with oral doxycycline. *Arthritis Rheum* 1992; 35:1150-1159
 141. Abatangelo G, Botti P, Del Bue M, et al: Intraarticular sodium hyaluronate injections in the Pond-Nuki experimental model of osteoarthritis in dogs—I. Biochemical results. *Clin Orthop* 1989; (241):278-285
 142. Schiavinato A, Lini E, Guidolin D, et al: Intraarticular sodium hyaluronate injections in the Pond-Nuki experimental model of osteoarthritis in dogs—II. Morphological findings. *Clin Orthop* 1989; (241):286-299
 143. Peyron JG: Intraarticular hyaluronan injections in the treatment of osteoarthritis: State-of-the-art review. *J Rheumatol Suppl* 1993; 39:10-15
 144. Dahlberg L, Lohmander LS, Ryd L: Intraarticular injections of hyaluronan in patients with cartilage abnormalities and knee pain—A one-year double-blind, placebo-controlled study. *Arthritis Rheum* 1994; 37:521-528
 145. McCarthy GM, McCarty DJ: Effect of topical capsaicin in the therapy of painful osteoarthritis of the hands. *J Rheumatol* 1992; 19:604-607
 146. Altman R, Aven A, Holmburg C, Pfeifer L, Sack M, Young G: Capsaicin cream 0.025% as monotherapy for osteoarthritis: A double-blind study. *Semin Arthritis Rheum* 1994; 23(suppl 3):25-33
 147. Bert JM: Role of abrasion arthroplasty and debridement in the management of osteoarthritis of the knee. *Rheum Dis Clin North Am* 1993; 19:725-739
 148. Ike RW, Arnold WJ, Rothschild EW, et al: Tidal irrigation versus conservative medical management in patients with osteoarthritis of the knee: A prospective randomized study—Tidal Irrigation Cooperating Group. *J Rheumatol* 1992; 19:772-779
 149. Brittberg M, Lindahl A, Nilsson A, Ohlsson C, Isaksson O, Peterson L: Treatment of deep cartilage defects in the knee with autologous chondrocyte transplantation. *N Engl J Med* 1994; 331:889-895

This article is one of a series on topics in primary care in which common diagnostic or therapeutic problems encountered in primary care practice are presented. Physicians interested in contributing to the series are encouraged to contact the series' editors.

STEPHEN J. McPHEE, MD
TERRIE MENDELSON, MD
Assistant Editors